

## Diltiazem Hydrochloride Extended-Release Capsules

<b>Type of Posting</b>	Revision Bulletin
<b>Posting Date</b>	31–Jan–2020
<b>Official Date</b>	01–Feb–2020
<b>Expert Committee</b>	Chemical Medicines Monographs 2
<b>Reason for Revision</b>	Compliance

In accordance with the Rules and Procedures of the 2015–2020 Council of Experts, the Chemical Medicines Monographs 2 Expert Committee has revised the Diltiazem Hydrochloride Extended-Release Capsules monograph. The purpose for the revision is to add *Dissolution Test 21* to accommodate FDA-approved drug products with different dissolution conditions and tolerances than the existing dissolution tests. The revision also necessitates a change in the table numbering in the test for *Organic Impurities*.

The Diltiazem Hydrochloride Extended-Release Capsules Revision Bulletin supersedes the currently official monograph.

Should you have any questions, please contact Josan Thomas, Scientific Liaison (+91-404-448-8948 or [josan.thomas@usp.org](mailto:josan.thomas@usp.org)).

## Diltiazem Hydrochloride Extended-Release Capsules

### DEFINITION

Diltiazem Hydrochloride Extended-Release Capsules contain NLT 90.0% and NMT 110.0% of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ).

### IDENTIFICATION

- **A.** The UV-Vis spectrum of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay*.
- **B.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay*.

### ASSAY

#### PROCEDURE

**Solution A:** 0.79 g/L of ammonium bicarbonate in water. Adjust with diluted ammonia solution or acetic acid to a pH of 8.0.

**Solution B:** Acetonitrile

**Mobile phase:** See *Table 1*.

**Table 1**

Time (min)	Solution A (%)	Solution B (%)
0	95	5
2.0	95	5
5.0	60	40
13.0	60	40
16.0	30	70
20.0	30	70
20.1	95	5
25.0	95	5

**Diluent:** Acetonitrile and water (40:60)

**Standard solution:** 0.05 mg/mL of USP Diltiazem Hydrochloride RS in *Diluent*

**Sample stock solution:** Nominally 0.5 mg/mL of diltiazem hydrochloride from the Capsules in *Diluent* prepared as follows. Transfer a portion of finely powdered contents of NLT 20 Capsules to a suitable volumetric flask. Add *Diluent* equivalent to 80% of the flask volume, mechanically shake for 30 min, and sonicate for 60 min. Dilute with the *Diluent* to volume. Centrifuge and use the supernatant.

**Sample solution:** Nominally 0.05 mg/mL of diltiazem hydrochloride prepared in *Diluent* from *Sample stock solution*

#### Chromatographic system

(See *Chromatography* <621>, *System Suitability*.)

**Mode:** LC

**Detector:** UV 240 nm. For *Identification A*, use a diode array detector in the range of 190–400 nm.

**Column:** 2.1-mm × 15-cm; 1.7-μm packing L1

**Flow rate:** 0.3 mL/min

**Injection volume:** 2.0 μL

#### System suitability

**Sample:** *Standard solution*

#### Suitability requirements

**Tailing factor:** NMT 2.0

**Relative standard deviation:** NMT 1.0%

#### Analysis

**Samples:** *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) in the portion of Capsules taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

$r_U$  = peak response of diltiazem from the *Sample solution*

$r_S$  = peak response of diltiazem from the *Standard solution*

$C_S$  = concentration of USP Diltiazem Hydrochloride RS in the *Standard solution* (mg/mL)

$C_U$  = nominal concentration of diltiazem hydrochloride in the *Sample solution* (mg/mL)

**Acceptance criteria:** 90.0%–110.0%

### PERFORMANCE TESTS

#### Change to read:

#### DISSOLUTION <711>

**For products labeled for dosing every 12 h**

**Test 1:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 1*.

**Medium:** Water; 900 mL

**Apparatus 2:** 100 rpm

**Times:** 3, 9, and 12 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 2*.

**Table 2**

Time (h)	Amount Dissolved (%)
3	10–25
9	45–85
12	NLT 70

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 4:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 4*.

**Medium:** Water; 900 mL

**Apparatus 2:** 100 rpm

**Times:** 4, 8, 12, and 24 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 3*.

**Table 3**

Time (h)	Amount Dissolved (%)
4	10–25
8	35–60
12	55–80

**Table 3** (continued)

Time (h)	Amount Dissolved (%)
24	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 5:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 5*.

**Medium:** 0.05 M phosphate buffer, pH 7.2; 900 mL

**Apparatus 2:** 50 rpm

**Times:** 1, 3, and 8 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 4*.

**Table 4**

Time (h)	Amount Dissolved (%)
1	NMT 15
3	45–70
8	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 10:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 10*.

**Buffer:** Dissolve 7.1 g of anhydrous dibasic sodium phosphate in 1000 mL of water, and adjust with phosphoric acid to a pH of 6.5.

**Medium:** *Buffer*; 900 mL

**Apparatus 1:** 100 rpm

**Times:** 1, 6, 9, and 24 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 5*.

**Table 5**

Time (h)	Amount Dissolved (%)
1	NMT 10
6	10–30
9	34–60
24	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**For products labeled for dosing every 24 h**

**Test 2:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 2*.

**Medium:** Water; 900 mL

**Apparatus 2:** 100 rpm

**Times:** 1, 4, 10, and 15 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 6*.

**Table 6**

Time (h)	Amount Dissolved (%)
1	5–20
4	30–50
10	70–90
15	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 3:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 3*.

**Medium:** 0.1 N hydrochloric acid; 900 mL

**Apparatus 2:** 100 rpm

**Times:** 6, 12, 18, 24, and 30 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 7*.

**Table 7**

Time (h)	Amount Dissolved (%)
6	20–45
12	25–50
18	35–70
24	NLT 70
30	NLT 85

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 6:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 6*.

**Medium:** Water; 900 mL

**Apparatus 1:** 100 rpm

**Times:** 2, 4, 8, 12, and 16 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 8*.

**Table 8**

Time (h)	Amount Dissolved (%)
2	NMT 25
4	25–50
8	60–85
12	NLT 70
16	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 7:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 7*.

**Buffer:** Transfer 115 mL of acetic acid to a 10-L volumetric flask, dilute with water to volume, and mix (*Solution A*). Transfer 165.4 g of anhydrous sodium acetate to a 10-L volumetric flask, dilute with water to volume, and mix (*Solution B*). Mix 4410 mL of *Solution A* with 1590 mL of *Solution B*. Adjust, if necessary, with the addition of *Solution A* or *Solution B* to a pH of  $4.2 \pm 0.05$ .

**Medium:** *Buffer*; 900 mL

**Apparatus 2:** 100 rpm

**Times:** 1, 4, 10, and 15 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 9*.

**Table 9**

Time (h)	Amount Dissolved (%)
1	NMT 10
4	15–35
10	65–85
15	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 8:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 8*.

**Medium:** Water; 900 mL

**Apparatus 2:** 100 rpm

**Times:** 1, 4, 10, and 15 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 10*.

**Table 10**

Time (h)	Amount Dissolved (%)
1	5–20

**Table 10** (continued)

Time (h)	Amount Dissolved (%)
4	30–50
10	60–90
15	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 9:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 9*.

[NOTE—Perform the test separately in each of the two media.]

**Medium 1:** 0.1 N hydrochloric acid; 900 mL

**Medium 2:** Simulated intestinal fluid TS, prepared without enzyme and adjusted to a pH of  $7.5 \pm 0.1$ ; 900 mL

**Apparatus 2:** 75 rpm

**Time for Medium 1:** 2 h

**Times for Medium 2:** 2, 12, 18, and 24 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 11*.

**Table 11**

Time (h)	Amount Dissolved, Medium 1 (%)	Amount Dissolved, Medium 2 (%)
2	0–5	20–45
12	—	35–55
18	—	NLT 60
24	—	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 11:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 11*.

**Medium:** 0.1 N hydrochloric acid; 900 mL

**Apparatus 2:** 100 rpm

**Times:** 1, 6, 12, and 18 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 12*.

**Table 12**

Time (h)	Amount Dissolved (%)
1	NMT 10
6	30–40
12	36–58
18	NLT 85

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 12:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 12*. Proceed as directed in *Dissolution* <711>, *Procedure, Apparatus 1 and Apparatus 2, Extended-Release Dosage Forms*.

**Medium:** Water; 900 mL

**Apparatus 1:** 100 rpm

**Times:** 2, 8, 14, and 24 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 13*.

**Table 13**

Time (h)	Amount Dissolved (%)
2	NMT 20
8	30–55
14	NLT 65
24	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 13:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 13*. Proceed as directed in *Dissolution* <711>, *Procedure, Apparatus 1 and Apparatus 2, Extended-Release Dosage Forms*.

**Medium:** Water; 900 mL

**Apparatus 1:** 100 rpm

**Times:** 2, 8, 14, and 24 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 14*.

**Table 14**

Time (h)	Amount Dissolved (%)
2	NMT 20
8	30–55
14	60–80
24	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 14:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 14*. Proceed as directed in *Dissolution* <711>, *Procedure, Apparatus 1 and Apparatus 2, Extended-Release Dosage Forms*.

**Medium:** 0.1 N hydrochloric acid; 900 mL

**Apparatus 2:** 100 rpm

**Times:** 6, 12, 18, 24, and 30 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 15*.

**Table 15**

Time (h)	Amount Dissolved (%)
6	20–45
12	25–50
18	35–70
24	NLT 70
30	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 15:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 15*. Proceed as directed in *Dissolution* <711>, *Procedure, Apparatus 1 and Apparatus 2, Extended-Release Dosage Forms*.

**Medium:** 0.05 M phosphate buffer, pH 7.5; 900 mL

**Apparatus 2:** 75 rpm

**Times:** 2, 4, 8, 12, and 16 h

**Detector:** UV 237 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Sample per *Dissolution* <711>. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Tolerances:** See *Table 16*.

**Table 16**

Time (h)	Amount Dissolved (%)
2	NMT 25
4	20–40
8	60–85
12	NLT 70
16	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 16:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 16*.

**Medium, Apparatus 2, Times, Standard solution, and**

**Sample solution:** Proceed as directed for *Test 3*.

**Detector:** UV 238 nm

**Tolerances:** See *Table 17*.

**Table 17**

Time (h)	Amount Dissolved (%)
6	20–45
12	30–55

**Table 17** (continued)

Time (h)	Amount Dissolved (%)
18	40–75
24	NLT 70
30	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 17:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 17*.

**Medium:** 0.1 N hydrochloric acid; 900 mL

**Apparatus 2:** 100 rpm, with wire helix sinkers

**Times:** 6, 12, and 30 h

**Detector:** UV 238 nm

**Standard solution:** USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** Dilute with *Medium*, if necessary, to a concentration that is similar to that of the *Standard solution*.

**Blank:** *Medium*

**Tolerances:** See *Table 18*.

**Table 18**

Time (h)	Amount Dissolved (%)
6	20–40
12	35–55
30	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 18:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 18*.

**Medium:** Water; 900 mL

**Apparatus 1:** 100 rpm

**Times:** 2, 4, 8, and 12 h

**Detector:** UV 237 nm

**Standard stock solution:** 0.28 mg/mL of USP Diltiazem Hydrochloride RS in *Medium* prepared as follows. To a suitable amount of USP Diltiazem Hydrochloride RS in a suitable volumetric flask, add *Medium* to 50% of the volume of the flask and sonicate for 5 min to dissolve. Dilute with *Medium* to volume.

**Standard solution:** 0.014 mg/mL of USP Diltiazem Hydrochloride RS in *Medium* from the *Standard stock solution*

**Sample solution:** At the times specified, withdraw 10 mL of the solution under test. Replace the aliquots withdrawn for analysis with equal volumes of fresh portions of *Medium* maintained at 37°. Pass the solution through a PVDF filter of 0.45- $\mu$ m pore size. Discard the first 2 mL of filtrate. Dilute with *Medium* to a concentration that is similar to that of the *Standard solution*.

**Analysis**

**Samples:** *Standard solution* and *Sample solution*

Calculate the concentration of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) in the sample withdrawn from the vessel at each time point *i*:

$$\text{Result} = (A_U/A_S) \times C_S \times D$$

$A_U$  = absorbance of diltiazem from the *Sample solution* at each time point

$A_S$  = absorbance of diltiazem from the *Standard solution*

$C_S$  = concentration of USP Diltiazem Hydrochloride RS in the *Standard solution* (mg/mL)

$D$  = dilution factor for the *Sample solution*

Calculate the percentage of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at each time point *i*:

$$\text{Result}_1 = C_i \times V \times (1/L) \times 100$$

$$\text{Result}_2 = [(C_2 \times V) + (C_1 \times V_3)] \times (1/L) \times 100$$

$$\text{Result}_3 = \{(C_3 \times V) + [(C_2 + C_1) \times V_3]\} \times (1/L) \times 100$$

$$\text{Result}_4 = \{(C_4 \times V) + [(C_3 + C_2 + C_1) \times V_3]\} \times (1/L) \times 100$$

$C_i$  = concentration of diltiazem hydrochloride in the *Sample solution* withdrawn at the specified time point (mg/mL)

$V$  = volume of *Medium*, 900 mL

$L$  = label claim (mg/Capsule)

$V_3$  = volume of the *Sample* withdrawn at each time point (mL)

**Tolerances:** See *Table 19*.

**Table 19**

Time Point (i)	Time (h)	Amount Dissolved (%)
1	2	NMT 20
2	4	33–58
3	8	68–88
4	12	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

**Test 19:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 19*.

**Medium:** 0.1 N hydrochloric acid; 900 mL

**Temperature:** 37.0°–37.5°

**Apparatus 2:** 100 rpm, with a suitable sinker

**Times:** 1, 4, 12, 18, and 24 h

**Detector:** UV 238 nm

**Cell:** 0.5 mm

**Standard solution:** 0.4 mg/mL of USP Diltiazem Hydrochloride RS in *Medium*

**Sample solution:** A portion of the solution under test at the time points specified

**Analysis**

**Samples:** *Standard solution* and *Sample solution*

Use an automatic dissolution system with an appropriate dissolution software.

Calculate the percentage of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at each time point *i*:

$$\text{Result} = (A_U/A_S) \times (C_S/L) \times V \times 100$$

$A_U$  = absorbance of diltiazem from the *Sample solution* at each time point

$A_S$  = absorbance of diltiazem from the *Standard solution*

## 6 Diltiazem

Revision Bulletin  
Official February 1, 2020

$C_s$  = concentration of USP Diltiazem Hydrochloride RS in the *Standard solution* (mg/mL)  
 $L$  = label claim (mg/Capsule)  
 $V$  = volume of *Medium*, 900 mL

**Tolerances:** See *Table 20*.

**Table 20**

Time Point (i)	Time (h)	Amount Dissolved (%)
1	1	NMT 10
2	4	15–35
3	12	30–50
4	18	50–70
5	24	NLT 85

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* (711), *Acceptance Table 2*.

**Test 20:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 20*.

*Dissolution Test 20* is suitable for products labeled to contain 360 mg of diltiazem hydrochloride.

**Medium:** 0.1 N hydrochloric acid; 900 mL

**Apparatus 2:** 100 rpm

**Times:** 6, 12, 18, and 24 h

**Detector:** UV 237 nm

**Cell:** 0.05 cm

**Standard solution:** 0.4 mg/mL of USP Diltiazem Hydrochloride RS prepared as follows. Transfer a suitable amount of USP Diltiazem Hydrochloride RS into a suitable volumetric flask, and add methanol to 5% of the total volume of the flask to dissolve. Dilute with *Medium* to volume.

**Sample solution:** Pass a portion of the solution under test through a suitable filter at the time points specified.

**Blank:** *Medium*

### Analysis

**Samples:** *Standard solution* and *Sample solution*

Calculate the concentration of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) in the sample withdrawn from the vessel at each time point (i):

$$\text{Result} = (A_U/A_S) \times C_S$$

$A_U$  = absorbance of diltiazem from the *Sample solution* at each time point

$A_S$  = absorbance of diltiazem from the *Standard solution*

$C_S$  = concentration of USP Diltiazem Hydrochloride RS in the *Standard solution* (mg/mL)

Calculate the percentage of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at each time point (i):

$$\text{Result}_1 = C_1 \times V \times (1/L) \times 100$$

$$\text{Result}_2 = \{[C_2 \times (V - V_S)] + (C_1 \times V_S)\} \times (1/L) \times 100$$

$$\text{Result}_3 = \{[C_3 \times [V - (2 \times V_S)]] + [(C_2 + C_1) \times V_S]\} \times (1/L) \times 100$$

$$\text{Result}_4 = \{[C_4 \times [V - (3 \times V_S)]] + [(C_3 + C_2 + C_1) \times V_S]\} \times (1/L) \times 100$$

$C_i$  = concentration of diltiazem hydrochloride in the *Sample solution* withdrawn at the specified time point (mg/mL)

$V$  = volume of *Medium*, 900 mL

$L$  = label claim (mg/Capsule)

$V_S$  = volume of the *Sample* withdrawn at each time point (mL)

**Tolerances:** See *Table 21*.

**Table 21**

Time Point (i)	Time (h)	Amount Dissolved (%)
1	6	30–50
2	12	35–55
3	18	50–70
4	24	NLT 85

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* (711), *Acceptance Table 2*.

**Test 21:** If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 21*.

**Medium:** 0.1 N hydrochloric acid; 900 mL, deaerated

**Apparatus 2:** 100 rpm

**Times:** 2, 4, 14, 18, and 24 h

**Standard stock solution:** 1.33 mg/mL of USP Diltiazem Hydrochloride RS in *Medium*

**Standard solution:** (L/900) mg/mL of USP Diltiazem Hydrochloride RS from the *Standard stock solution* in *Medium*, where L is the label claim in mg/Capsule

**Sample solution:** Pass a portion of the solution under test at the time points specified through a suitable filter.

### Instrumental conditions

(See *Ultraviolet-Visible Spectroscopy* (857).)

**Mode:** UV

**Analytical wavelength:** 237 nm for 120 mg, 180 mg, and 240 mg strength capsules.

260 nm for 300 mg and 360 mg strength capsules.

**Cell:** 1 mm for 120 mg, 180 mg, and 240 mg strength capsules.

2 mm for 300 mg and 360 mg strength capsules.

**Blank:** *Medium*

### Analysis

**Samples:** *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at each time point (i):

$$\text{Result} = (A_U/A_S) \times (C_S/L) \times V \times 100$$

$A_U$  = absorbance of diltiazem from the *Sample solution* at each time point

$A_S$  = absorbance of diltiazem from the *Standard solution*

$C_S$  = concentration of USP Diltiazem Hydrochloride RS in the *Standard solution* (mg/mL)

$L$  = label claim (mg/Capsule)

$V$  = volume of *Medium*, 900 mL

**Tolerances:** See *Table 22*.

**Table 22**

Time Point (j)	Time (h)	Amount Dissolved (%)
1	2	NMT 20
2	4	25–45
3	14	35–55
4	18	70–90
5	24	NLT 80

The percentages of the labeled amount of diltiazem hydrochloride ( $C_{22}H_{26}N_2O_4S \cdot HCl$ ) dissolved at the times specified conform to *Dissolution* (711), *Acceptance Table 2*. (RB 1-Feb-2020)

- **UNIFORMITY OF DOSAGE UNITS** (905): Meet the requirements

**IMPURITIES**

**Change to read:**

• **ORGANIC IMPURITIES**

**Solution A, Solution B, Mobile phase, Diluent, and Chromatographic system:** Proceed as directed in the Assay.

**Standard solution:** 2.5 µg/mL each of USP Desacetyl Diltiazem Hydrochloride RS and USP Diltiazem Hydrochloride RS in *Diluent*

**Sample solution:** Nominally 0.5 mg/mL of diltiazem hydrochloride from the Capsules in *Diluent* prepared as follows. Transfer a portion of the finely powdered contents of NLT 20 Capsules to a suitable volumetric flask. Add *Diluent* equivalent to 80% of the flask volume, mechanically shake for 30 min, and sonicate for 60 min. Dilute with *Diluent* to volume. Centrifuge and use the supernatant.

**System suitability**

**Sample:** *Standard solution*

[NOTE—For relative retention times see **Table 23**. (RB 1-Feb-2020)]

**Suitability requirements**

**Resolution:** NLT 2.0 between desacetyl diltiazem and diltiazem

**Relative standard deviation:** NMT 3.0%

**Analysis**

**Samples:** *Standard solution* and *Sample solution*

Calculate the percentage of desacetyl diltiazem hydrochloride in the portion of Capsules taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

$r_U$  = peak response of desacetyl diltiazem from the *Sample solution*

$r_S$  = peak response of desacetyl diltiazem from the *Standard solution*

$C_S$  = concentration of USP Desacetyl Diltiazem Hydrochloride RS in the *Standard solution* (µg/mL)

$C_U$  = nominal concentration of diltiazem hydrochloride in the *Sample solution* (µg/mL)

Calculate the percentage of any individual unspecified impurity in the portion of Capsules taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

$r_U$  = peak response of each unspecified impurity from the *Sample solution*

$r_S$  = peak response of diltiazem from the *Standard solution*

$C_S$  = concentration of USP Diltiazem Hydrochloride RS in the *Standard solution* (µg/mL)

$C_U$  = nominal concentration of diltiazem hydrochloride in the *Sample solution* (µg/mL)

**Acceptance criteria:** See **Table 23**. (RB 1-Feb-2020) Disregard limit: 0.05%.

**Table 23** (RB 1-Feb-2020)

Name	Relative Retention Time	Acceptance Criteria, NMT (%)
Diltiazem related compound H <sup>a, b</sup>	0.44	—
Diltiazem related compound G <sup>b, c</sup>	0.52	—
Diltiazem related compound C <sup>b, d</sup>	0.58	—
Diltiazem related compound D <sup>b, e</sup>	0.61	—
Diltiazem related compound E <sup>b, f</sup>	0.66	—
Desacetyl diltiazem <sup>g</sup>	0.75	1.5
Diltiazem related compound A <sup>b, h</sup>	0.83	—
Diltiazem related compound B <sup>b, i</sup>	0.89	—
Diltiazem	1.0	—
Any individual unspecified impurity	—	0.2
Total impurities	—	2.0

<sup>a</sup> (2S,3S)-5-(2-Aminoethyl)-3-hydroxy-2-(4-hydroxyphenyl)-2,3-dihydro-1,5-benzothiazepine-4(5H)-one.

<sup>b</sup> These are impurities related to the drug substance. They are not to be reported for the drug product and should not be included in the total impurities.

<sup>c</sup> (2S,3S)-3-Hydroxy-2-(3-methoxyphenyl)-5-(2-(methylamino)ethyl)-2,3-dihydrobenzo[b][1,4]thiazepin-4(5H)-one.

<sup>d</sup> (2S,3S)-5-[2-(Dimethylamino)ethyl]-2-(4-hydroxyphenyl)-4-oxo-2,3,4,5-tetrahydro-1,5-benzothiazepin-3-yl acetate.

<sup>e</sup> (2S,3S)-2-(4-Methoxyphenyl)-5-[2-(methylamino)ethyl]-4-oxo-2,3,4,5-tetrahydro-1,5-benzothiazepin-3-yl acetate.

<sup>f</sup> (2S,3S)-3-Hydroxy-2-(4-methoxyphenyl)-2,3-dihydro-1,5-benzothiazepin-4(5H)-one.

<sup>g</sup> *d-cis*-3-Hydroxy-2,3-dihydro-5-[2-(dimethylamino)ethyl]-2-(*p*-methoxyphenyl)-1,5-benzothiazepin-4(5H)-one. The acceptance criteria for this impurity is based on the hydrochloride form.

<sup>h</sup> (2R,3S)-5-[2-(Dimethylamino)ethyl]-2-(4-methoxyphenyl)-4-oxo-2,3,4,5-tetrahydro-1,5-benzothiazepin-3-yl acetate.

<sup>i</sup> (2S,3S)-2-(4-Methoxyphenyl)-4-oxo-2,3,4,5-tetrahydro-1,5-benzothiazepin-3-yl acetate.

**ADDITIONAL REQUIREMENTS**

- **PACKAGING AND STORAGE:** Preserve in tight containers. Store at controlled room temperature.

- **LABELING:** The labeling indicates the *Dissolution* test with which the product complies.

- **USP REFERENCE STANDARDS** (11)

USP Desacetyl Diltiazem Hydrochloride RS  
*d-cis*-3-Hydroxy-2,3-dihydro-5-[2-(dimethylamino)ethyl]-2-(*p*-methoxyphenyl)-1,5-benzothiazepin-4(5H)-one hydrochloride.  
 $C_{20}H_{24}N_2O_3S \cdot HCl$  408.95  
USP Diltiazem Hydrochloride RS