

## Potassium Chloride Extended-Release Tablets

<b>Type of Posting</b>	Revision Bulletin
<b>Posting Date</b>	31–Aug–2018
<b>Official Date</b>	01–Sep–2018
<b>Expert Committee</b>	Chemical Medicines Monographs 5
<b>Reason for Revision</b>	Compliance

In accordance with the Rules and Procedures of the 2015–2020 Council of Experts, the Chemical Medicines Monographs 5 Expert Committee has revised the Potassium Chloride Extended-Release Tablets monograph. The purpose for the revision is to add *Dissolution Test 2* to accommodate FDA-approved drug products with different tolerances than the existing dissolution test. *Labeling* information has been incorporated to support the inclusion of *Dissolution Test 2*.

The Potassium Chloride Extended-Release Tablets Revision Bulletin supersedes the currently official monograph.

Should you have any questions, please contact Ren-Hwa Yeh, Ph.D., Senior Scientific Liaison (301-998-6818 or [rhy@usp.org](mailto:rhy@usp.org)).

## Potassium Chloride Extended-Release Tablets

### DEFINITION

Potassium Chloride Extended-Release Tablets contain NLT 90.0% and NMT 110.0% of the labeled amount of potassium chloride (KCl).

### IDENTIFICATION

- **A. IDENTIFICATION TESTS—GENERAL** (191), *Chemical Identification Tests, Potassium*

**Sample solution:** A portion of the filtrate, obtained as directed for the designated *Sample stock solution* in the Assay

**Acceptance criteria:** Meet the requirements

- **B. IDENTIFICATION TESTS—GENERAL** (191), *Chemical Identification Tests, Chloride*

**Sample solution:** A portion of the filtrate, obtained as directed for the designated *Sample stock solution* in the Assay

**Acceptance criteria:** Meet the requirements

### ASSAY

#### PROCEDURE

[NOTE—If necessary, first score nonsugar-coated Tablets. Retain a portion of the filtrate of either *Sample stock solution 1* or *Sample stock solution 2* for use in *Identification A* and *B*.]

**Standard stock solution:** 19.07 µg/mL of potassium chloride, previously dried at 105° for 2 h, in water. This solution contains 10 µg/mL of potassium.

**Standard solutions:** To separate 100-mL volumetric flasks transfer 10.0, 15.0, and 20.0 mL, respectively, of *Standard stock solution*. To each flask add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume. The *Standard solutions* contain 1.0, 1.5, and 2.0 µg/mL of potassium, respectively.

#### Sample preparation 1

**Sample stock solution 1:** Nominally 0.06 mg/mL of potassium chloride prepared as follows. Place NLT 20 Tablets in a suitable container with 400 mL of water, heat to boiling, and boil for 20 min. Allow to cool, transfer the solution to a 1000-mL volumetric flask, and dilute with water to volume. Filter and discard the first 20 mL of the filtrate. Transfer a measured volume of the subsequent filtrate, equivalent to 60 mg of potassium chloride, to a 1000-mL volumetric flask, and dilute with water to volume.

**Sample solution 1:** Nominally 3 µg/mL of potassium chloride prepared as follows. Transfer 5.0 mL of *Sample stock solution 1* to a 100-mL volumetric flask, add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume.

**Sample preparation 2** (for formulations containing crystals coated with hydrophobic polymers)

**Sample stock solution 2:** Nominally 0.06 mg/mL of potassium chloride prepared as follows. Place NLT 20 Tablets in a 2000-mL volumetric flask. Add 1200 mL of a mixture of acetonitrile and water (1:1), and shake by mechanical means, or stir using a magnetic bar for 90 min. Dilute with the mixture of acetonitrile and water (1:1) to volume. Allow to stand for 90 min. Pass through a filter of 0.2-µm pore size. Transfer a measured volume of the filtrate, and quantitatively dilute with water to obtain a solution with a concentration of 0.06 mg/mL.

[NOTE—Alternatively, *Sample stock solution 2* can be prepared by the following procedure. Nominally 0.15 mg/mL of potassium chloride from NLT 20 finely

powdered Tablets, prepared as follows. Transfer an appropriate amount of the powder, equivalent to about 5–6 Tablets, to a suitable volumetric flask, add 10% of the final flask volume of acetone, and sonicate for 45 min with intermittent shaking. Add 80% of the final flask volume of water and sonicate for 45 min with intermittent shaking. Cool to room temperature and dilute with water to volume. Centrifuge a portion of the solution at 5000 rpm for 10 min. Transfer an appropriate amount of the supernatant to a 100-mL volumetric flask and dilute with water to volume to obtain a solution with a concentration of 0.15 mg/mL.]

**Sample solution 2:** Nominally 3 µg/mL of potassium chloride prepared as follows. Transfer an appropriate amount of *Sample stock solution 2* to a 100-mL volumetric flask, add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume.

#### Instrumental conditions

(See *Atomic Absorption Spectroscopy* (852).)

**Mode:** Atomic absorption spectrophotometry

**Analytical wavelength:** Potassium emission line at 766.5 nm

**Lamp:** Potassium hollow-cathode

**Flame:** Air–acetylene

**Blank:** Water

#### Analysis

**Samples:** *Standard solutions*, *Sample solution 1* or *Sample solution 2*, and *Blank*

Plot the absorbances of the *Standard solutions* versus the concentration of potassium, in µg/mL, and draw the straight line best fitting the three plotted points. From the graph, determine the concentration of potassium in the *Sample solution* (µg/mL).

Calculate the percentage of the labeled amount of potassium chloride (KCl) in each Tablet taken:

$$\text{Result} = (C/C_U) \times (M_r/A_r) \times 100$$

C	= concentration of potassium in the <i>Sample solution</i> as determined in this test (µg/mL)
C <sub>U</sub>	= nominal concentration of potassium chloride in the <i>Sample solution</i> (µg/mL)
M <sub>r</sub>	= molecular weight of potassium chloride, 74.55
A <sub>r</sub>	= atomic weight of potassium, 39.10

**Acceptance criteria:** 90.0%–110.0%

### PERFORMANCE TESTS

#### Change to read:

#### DISSOLUTION (711)

▲**Test 1**▲ (RB 1-Sep-2018)

**Medium:** Water; 900 mL

**Apparatus 2:** 50 rpm

**Time:** 2 h

**Standard stock solution:** 19.07 µg/mL of potassium chloride, previously dried at 105° for 2 h, in water. This solution contains 10 µg/mL of potassium.

**Standard solutions:** To separate 100-mL volumetric flasks transfer 10.0, 15.0, and 20.0 mL, respectively, of *Standard stock solution*. To each flask add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume. The *Standard solutions* contain, respectively, 1.0, 1.5, and 2.0 µg/mL of potassium.

**Sample stock solution:** Filter the solution under test, and dilute with *Medium* to obtain a solution containing nominally 60 µg/mL of potassium chloride.

**Sample solution:** Transfer 5.0 mL of the *Sample stock solution* to a 100-mL volumetric flask. Add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume.

#### Instrumental conditions

(See *Atomic Absorption Spectroscopy* (852).)

**Mode:** Atomic absorption spectrophotometry

**Analytical wavelength:** Potassium emission line at 766.5 nm

**Lamp:** Potassium hollow-cathode

**Flame:** Air-acetylene

**Blank:** Water

#### Analysis

**Samples:** *Standard solutions*, *Sample solution*, and *Blank*  
Plot the absorbances of the *Standard solutions* versus the concentration of potassium, in µg/mL, and draw the straight line best fitting the three plotted points. From the graph, determine the concentration of potassium in the *Sample solution* (µg/mL).

Calculate the percentage of the labeled amount of potassium chloride (KCl) dissolved:

$$\text{Result} = [C \times D \times (V/L)] \times (M_r/A_r) \times 100$$

- C* = concentration of potassium in the *Sample solution* as determined in this test (µg/mL)  
*D* = dilution factor of the *Sample solution*  
*V* = volume of *Medium*, 900 mL  
*L* = labeled amount of potassium chloride (µg/ Tablet)  
*M<sub>r</sub>* = molecular weight of potassium chloride, 74.55  
*A<sub>r</sub>* = atomic weight of potassium, 39.10

**Tolerances:** NMT 35% (*Q*) of the labeled amount of potassium chloride (KCl) is dissolved in 2 h. The requirements are met if the quantities dissolved from the Tablets tested conform to *Table 1* instead of the table shown in *Dissolution* (711).

**Table 1**

Stage	Number Tested	Acceptance Criteria
<i>S</i> <sub>1</sub>	6	Each unit is within the range $Q \pm 30\%$ .
<i>S</i> <sub>2</sub>	6	Average of 12 units ( <i>S</i> <sub>1</sub> + <i>S</i> <sub>2</sub> ) is within the range between $Q - 30\%$ and $Q + 35\%$ , and no unit is outside the range $Q \pm 40\%$ .
<i>S</i> <sub>3</sub>	12	Average of 24 units ( <i>S</i> <sub>1</sub> + <i>S</i> <sub>2</sub> + <i>S</i> <sub>3</sub> ) is within the range between $Q - 30\%$ and $Q + 35\%$ , and NMT 2 units are outside the range $Q \pm 40\%$ .

▲**Test 2:** If the product complies with this procedure, the labeling indicates that it meets USP *Dissolution Test 2*.

**Standard stock solution and Standard solutions:**

Prepare as directed in *Test 1*.

**Medium:** Water; 900 mL

**Apparatus 2:** 50 rpm

**Times:** 1, 2, 4, and 8 h

**Sample stock solution:** Transfer 4.0 mL of the solution under test into either a 50-mL volumetric flask (for 750-mg Tablet) or a 100-mL volumetric flask (for 1500-mg Tablet), dilute with water to volume, and filter.

**Sample solution:** Transfer 4.0 mL of the *Sample stock solution* to a 100-mL volumetric flask. Add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume.

**Blank solution:** To a 100-mL volumetric flask, add 2.0 mL of sodium chloride solution (1 in 5) and 1.0 mL of hydrochloric acid, and dilute with water to volume.

**Instrumental conditions:** Proceed as directed in *Test 1*, except do not use the *Blank*.

#### System suitability

**Samples:** *Standard solutions*

#### Suitability requirements

**Linearity:** Correlation coefficient NLT 0.99

**Relative standard deviation:** NMT 5.0% from 5

replicate analyses of the 1.5-µg/mL *Standard solution*

#### Analysis

**Samples:** 1.5-µg/mL *Standard solution*, *Sample solution*, and *Blank solution*

Calculate the percentage of the labeled amount of potassium chloride (KCl) dissolved:

$$\text{Result}_i = [(A_U/A_S) \times C_S \times D \times (V/L)] \times (M_r/A_r) \times 100$$

*A<sub>U</sub>* = absorbance of potassium in the *Sample solution*

*A<sub>S</sub>* = absorbance of potassium in the *Standard solution*

*C<sub>S</sub>* = concentration of potassium in the *Standard solution* (µg/mL)

*D* = dilution factor of the *Sample solution*

*V* = volume of *Medium*, 900 mL

*L* = labeled amount of potassium chloride (µg/ Tablet)

*M<sub>r</sub>* = molecular weight of potassium chloride, 74.55

*A<sub>r</sub>* = atomic weight of potassium, 39.10

**Tolerances:** See *Table 2*.

**Table 2**

Time Point ( <i>i</i> )	Time (h)	Amount Dissolved (%)	
		750 mg/Tablet	1500 mg/Tablet
1	1	10–30	5–25
2	2	30–50	25–45
3	4	60–80	55–75
4	8	NLT 80	NLT 85

The percentages of the labeled amount of potassium chloride (KCl), dissolved at the times specified, conform to *Dissolution* (711), *Acceptance Table 2*. ▲ (RB 1-Sep-2018)

• **UNIFORMITY OF DOSAGE UNITS** (905): Meet the requirements

#### ADDITIONAL REQUIREMENTS

• **PACKAGING AND STORAGE:** Preserve in tight containers, and store at a temperature not exceeding 30°.

#### Change to read:

• **LABELING:** The label states with which *Sample preparation* in the *Assay* the product complies only if *Sample preparation 1* is not used. ▲When more than one *Dissolution* test is given, the labeling states the *Dissolution* test used only if *Test 1* is not used. ▲ (RB 1-Sep-2018)