

Quetiapine Extended-Release Tablets

Type of Posting	Revision Bulletin
Posting Date	29–Mar–2019
Official Date	01–Apr–2019
Expert Committee	Chemical Medicines Monographs 4
Reason for Revision	Compliance

In accordance with the Rules and Procedures of the 2015–2020 Council of Experts, the Chemical Medicines Monographs 4 Expert Committee has revised the Quetiapine Extended-Release Tablets monograph. The purpose for the revision is to add *Dissolution Test 7* to accommodate FDA-approved drug products with different dissolution conditions and/or tolerances than the existing dissolution tests.

Existing references to reagents have been updated for consistency with the reagent entry names. For additional information about reagent cross-references, please see the related [Compendial Notice](#).

The revision also necessitates a change in the table numbering in the *Assay* and in the test for *Organic Impurities*.

The Quetiapine Extended-Release Tablets Revision Bulletin supersedes the currently official monograph.

Should you have any questions, please contact Claire Chisolm, Scientific Liaison (301-230-3215 or cnc@usp.org).

Quetiapine Extended-Release Tablets

DEFINITION

Quetiapine Extended-Release Tablets contain quetiapine fumarate [(C₂₁H₂₅N₃O₂S)₂ · C₄H₄O₄] equivalent to NLT 90.0% and NMT 110.0% of the labeled amount of quetiapine (C₂₁H₂₅N₃O₂S).

IDENTIFICATION

• A. INFRARED ABSORPTION <197F>

Standard solution: Transfer 10 mg of USP Quetiapine Fumarate RS to a suitable vial. Add 10 mL of acetone and cap the vial. Sonicate for about 10 min. Allow the solution to equilibrate to room temperature. Evaporate the acetone completely. Add 2 mL of chloroform. Gently swirl for several minutes. Pass through a suitable filter of 0.45-µm pore size. Use the filtrate.

Sample solution: Grind NLT 10 Tablets. Transfer an amount of powder equivalent to NLT 10 mg of quetiapine fumarate to a suitable vial. Add 10 mL of acetone and cap the vial. Sonicate for about 10 min. Allow the solution to equilibrate to room temperature. Evaporate the acetone completely. Add 2 mL of chloroform. Gently swirl for several minutes. Pass through a suitable filter of 0.45-µm pore size. Use the filtrate.

Acceptance criteria: Meet the requirements

- **B.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay*.

ASSAY

Change to read:

• PROCEDURE

Buffer: Dissolve 2.6 g/L of dibasic ammonium phosphate in water.

Mobile phase: Methanol, acetonitrile, and *Buffer* (54:7:39)

Diluent: Acetonitrile and water (50:50)

System suitability stock solution: 0.05 mg/mL of USP Quetiapine Related Compound H RS in *Mobile phase*

System suitability solution: 0.005 mg/mL of USP Quetiapine Related Compound H RS and 0.5 mg/mL of USP Quetiapine System Suitability RS in *Mobile phase* prepared as follows. Transfer 5 mg of USP Quetiapine System Suitability RS to a 10-mL volumetric flask. Add 7 mL of *Mobile phase* and sonicate to dissolve. Transfer 1 mL of *System suitability stock solution* to the volumetric flask. Dilute with *Mobile phase* to volume.

Standard solution: 0.2 mg/mL of USP Quetiapine Fumarate RS in *Mobile phase*

Sample stock solution: Transfer NLT 5 Tablets to a homogenizer vessel. Add 50 mL of acetonitrile, swirl to wet, and allow to stand for approximately 10 min. Add an additional 160 mL of *Diluent* and extract for about 10 min. Transfer the contents of the homogenizer to a 500-mL volumetric flask. Dilute with *Diluent* to volume. Pass a portion of the solution through a suitable filter of 0.45-µm pore size and use the filtrate.

Sample solution: Nominally 0.16–0.18 mg/mL of quetiapine from the *Sample stock solution* in *Mobile phase*

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: UV 230 nm

Column: 4.6-mm × 25-cm; 5-µm packing L7

Flow rate: 1.3 mL/min

Injection volume: 30 µL

Run time: NLT 2.5 times the retention time of quetiapine

System suitability

Samples: *System suitability solution* and *Standard solution*
[NOTE—See **▲ Table 8▲** (RB 1-Apr-2019) for the relative retention times.]

Suitability requirements

Resolution: NLT 1.5 between quetiapine related compound G and quetiapine related compound H; NLT 2.0 between the quetiapine desthoxy and quetiapine peaks; *System suitability solution*

Tailing factor: NMT 1.5, *Standard solution*

Relative standard deviation: NMT 2.0%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*
Calculate the percentage of the labeled amount of quetiapine (C₂₁H₂₅N₃O₂S) in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (M_{r1}/M_{r2}) \times N \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of USP Quetiapine Fumarate RS in the *Standard solution* (mg/mL)

C_U = nominal concentration of quetiapine in the *Sample solution* (mg/mL)

M_{r1} = molecular weight of quetiapine free base, 383.51

M_{r2} = molecular weight of quetiapine fumarate, 883.09

N = number of moles of quetiapine free base per mole of quetiapine fumarate, 2

Acceptance criteria: 90.0%–110.0%

PERFORMANCE TESTS

Change to read:

• DISSOLUTION <711>

Test 1

Medium 1: Citrate buffer, pH 4.8. Dissolve 9.6 g of anhydrous citric acid in 600 mL of water. Add 90 mL of 1 N sodium hydroxide. Dilute with water to 1 L; 900 mL.

Medium 2: Dissolve 17.9 g of dibasic sodium phosphate dodecahydrate in 400 mL of water. Add 460 mL of **▲1 N sodium hydroxide VS▲** (RB 1-Apr-2019) and dilute with water to 1 L; 100 mL.

[NOTE—It is recommended to check the pH of the mixture of 90 mL of *Medium 1* and 10 mL of *Medium 2*, which should be between 6.4 and 6.8. If the pH of the mixture is less than 6.4, 10 mL/L of **▲1 N sodium hydroxide VS▲** (RB 1-Apr-2019) may be added to *Medium 2*. If the pH of the mixture is greater than 6.8, 10 mL/L of **▲1 N hydrochloric acid VS▲** (RB 1-Apr-2019) may be added to *Medium 2*.]

Start the test with 900 mL of *Medium 1*. Add 100 mL of *Medium 2* to the vessel after 5 h of the test and continue the test.

Apparatus 1: 200 rpm

Times: 1, 6, 12, and 20 h

Diluent: *Medium 1* and *Medium 2* (90:10)

Standard solution: (L/400) mg/mL of USP Quetiapine Fumarate RS in *Diluent*, where L is the label claim in mg/ Tablet

Sample solution: Pass a suitable portion of the solution under test through a suitable filter.

Instrumental conditions

Mode: UV

Analytical wavelength: About 290 nm

Blank: Diluent

Analysis

Samples: Standard solution and Sample solution

Calculate the concentration, C_i , of quetiapine ($C_{21}H_{25}N_3O_2S$) in Medium (mg/mL) after time point (i):

$$C_i = (A_U/A_S) \times C_S \times (M_{r1}/M_{r2}) \times N$$

- A_U = absorbance of the Sample solution
 A_S = absorbance of the Standard solution
 C_S = concentration of quetiapine fumarate in the Standard solution (mg/mL)
 M_{r1} = molecular weight of quetiapine free base, 383.51
 M_{r2} = molecular weight of quetiapine fumarate, 883.09
 N = number of moles of quetiapine free base per mole of quetiapine fumarate, 2

Calculate the percentage of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at each time point (i):

$$\text{Result}_1 = C_1 \times V \times (1/L) \times 100$$

$$\text{Result}_2 = [(C_2 \times V) + (C_1 \times V_5)] \times (1/L) \times 100$$

$$\text{Result}_3 = \{(C_3 \times V) + [(C_2 + C_1) \times V_5]\} \times (1/L) \times 100$$

$$\text{Result}_4 = \{(C_4 \times V) + [(C_3 + C_2 + C_1) \times V_5]\} \times (1/L) \times 100$$

- C_i = concentration of quetiapine in Medium in the portion of sample withdrawn at each time point (mg/mL)
 V = volume of Medium, 900 mL for 1 h; 1000 mL for 6-, 12-, and 20-h time points
 L = label claim (mg/Tablet)
 V_5 = volume of the Sample solution withdrawn from the vessel and replaced with Medium (mL)

Tolerances: See Table 1.

Table 1

Time Point (i)	Time (h)	Amount Dissolved (%)
1	1	NMT 20
2	6	47–69
3	12	65–95
4	20	NLT 85

The percentages of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

Test 2: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 2*.

Medium: Water; 900 mL

Apparatus 1: 100 rpm

Times: 2, 4, 8, and 24 h

Standard solution: 0.03 mg/mL of USP Quetiapine Fumarate RS in water

Sample solution: Pass a suitable portion of the solution under test through a suitable filter of 0.45- μ m pore size. Discard the first few milliliters of filtrate. Replace the volume withdrawn with an equal volume of Medium. Dilute with Medium to a concentration that is similar to that of the Standard solution.

Instrumental conditions

Mode: UV

Analytical wavelength: 290 nm

Blank: Water

Analysis

Samples: Standard solution and Sample solution

Calculate the concentration, C_i , of quetiapine ($C_{21}H_{25}N_3O_2S$) in Medium (mg/mL) after each time point (i):

$$C_i = (A_U/A_S) \times C_S \times D \times (M_{r1}/M_{r2}) \times N$$

- A_U = absorbance of the Sample solution
 A_S = absorbance of the Standard solution
 C_S = concentration of quetiapine fumarate in the Standard solution (mg/mL)
 D = dilution factor for the Sample solution, if needed
 M_{r1} = molecular weight of quetiapine free base, 383.51
 M_{r2} = molecular weight of quetiapine fumarate, 883.09
 N = number of moles of quetiapine free base per mole of quetiapine fumarate, 2

Calculate the percentage of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at each time point (i):

$$\text{Result}_1 = C_1 \times V \times (1/L) \times 100$$

$$\text{Result}_2 = [(C_2 \times V) + (C_1 \times V_5)] \times (1/L) \times 100$$

$$\text{Result}_3 = \{(C_3 \times V) + [(C_2 + C_1) \times V_5]\} \times (1/L) \times 100$$

$$\text{Result}_4 = \{(C_4 \times V) + [(C_3 + C_2 + C_1) \times V_5]\} \times (1/L) \times 100$$

- C_i = concentration of quetiapine in Medium in the portion of sample withdrawn at each time point (mg/mL)
 V = volume of Medium, 900 mL
 L = label claim (mg/Tablet)
 V_5 = volume of the Sample solution withdrawn from the vessel and replaced with Medium (mL)

Tolerances: See Table 2.

Table 2

Time Point (i)	Time (h)	Amount Dissolved (%)
1	2	5–25
2	4	20–45
3	8	45–75
4	24	NLT 85

The percentages of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

Test 3: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 3*.

Medium: ▲0.1 N hydrochloric acid VS; ▲ (RB 1-Apr-2019) 900 mL

Apparatus 2: 50 rpm

Times: 1, 4, and 8 h

Standard solution: USP Quetiapine Fumarate RS, equivalent to (L/900) mg/mL of quetiapine in Medium, where L is the label claim in mg/Tablet

Sample solution: Pass a suitable portion of the solution under test through a suitable full flow filter of 10- μ m pore size.

Instrumental conditions

Mode: UV

Analytical wavelength: 295 nm

Cell

For 50-mg Tablets: 10 mm

For 150-, 200-, 300-, and 400-mg Tablets: 1 mm

Blank: Medium

Analysis

Samples: Standard solution and Sample solution

Calculate the percentage of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at each time point (i):

$$\text{Result} = (A_U/A_S) \times C_S \times V \times (1/L) \times (M_{r1}/M_{r2}) \times N \times 100$$

A_U = absorbance of the Sample solution

A_S = absorbance of the Standard solution

C_S = concentration of quetiapine fumarate in the Standard solution (mg/mL)

V = volume of Medium, 900 mL

L = label claim (mg/Tablet)

M_{r1} = molecular weight of quetiapine free base, 383.51

M_{r2} = molecular weight of quetiapine fumarate, 883.09

N = number of moles of quetiapine free base per mole of quetiapine fumarate, 2

Tolerances: See Table 3.

Table 3

Time Point (i)	Time (h)	Amount Dissolved (for 50-, 150-, and 200-mg Tablets) (%)	Amount Dissolved (for 300- and 400-mg Tablets) (%)
1	1	NMT 40	NMT 35
2	4	55–75	45–65
3	8	NLT 85	NLT 80

The percentages of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

Test 4: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 4*.

Medium: Water; 900 mL

Apparatus 2: 100 rpm

Times: 1, 4, 8, and 16 h

Standard solution: USP Quetiapine Fumarate RS, equivalent to ($L/900$) mg/mL of quetiapine in Medium, where L is the label claim in mg/Tablet

Sample solution: Pass a suitable portion of the solution under test through a suitable filter.

Instrumental conditions

Mode: UV

Analytical wavelength: 250 nm

Blank: Medium

Analysis

Samples: Standard solution and Sample solution

Calculate the percentage of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at each time point (i):

$$\text{Result} = (A_U/A_S) \times C_S \times V \times (1/L) \times (M_{r1}/M_{r2}) \times N \times 100$$

A_U = absorbance of the Sample solution

A_S = absorbance of the Standard solution

C_S = concentration of quetiapine fumarate in the Standard solution (mg/mL)

V = volume of Medium, 900 mL

L = label claim (mg/Tablet)

M_{r1} = molecular weight of quetiapine free base, 383.51

M_{r2} = molecular weight of quetiapine fumarate, 883.09

N = number of moles of quetiapine free base per mole of quetiapine fumarate, 2

Tolerances: See Table 4. The percentages of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

Test 5: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 5*.

Medium, Apparatus 1, Times, Standard solution,

Sample solution, Instrumental conditions, and

Analysis: Proceed as directed in *Dissolution Test 2*.

Tolerances: See Table 5.

Table 5

Time Point (i)	Time (h)	Amount Dissolved (%)
1	2	10–30
2	4	30–50
3	8	60–80
4	24	NLT 85

Table 4

Time Point (i)	Time (h)	Amount Dissolved (for 50-mg Tablets) (%)	Amount Dissolved (for 150-mg Tablets) (%)	Amount Dissolved (for 200-mg Tablets) (%)	Amount Dissolved (for 300-mg Tablets) (%)	Amount Dissolved (for 400-mg Tablets) (%)
1	1	NMT 20	NMT 20	NMT 20	NMT 15	NMT 15
2	4	30–55	35–55	28–48	22–42	22–42
3	8	60–85	65–90	60–85	52–76	50–75
4	16	NLT 85	NLT 85	NLT 85	NLT 85	NLT 85

The percentages of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

Test 6: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 6*.

Medium: pH 6.8 phosphate buffer (6.8 g/L of monobasic potassium phosphate and 0.9 g/L of sodium hydroxide in water. Adjust with 1 N sodium hydroxide VS or phosphoric acid to a pH of 6.8, and sonicate for NLT 10 min); 900 mL

Apparatus 2: 100 rpm, with sinker

Times: 1, 4, 8, and 16 h

Mobile phase: Methanol, trifluoroacetic acid, and water (40:0.1:60)

Standard solution: 0.1 mg/mL of USP Quetiapine Fumarate RS prepared as follows. Transfer an appropriate amount of USP Quetiapine Fumarate RS to a suitable volumetric flask, and add 5% of the final flask volume of methanol. Sonicate to dissolve, then dilute with *Medium* to volume.

Sample solution: Pass a suitable portion of the solution under test through a suitable filter of 0.45- μ m pore size. Discard the first few milliliters of filtrate.

Chromatographic system

(See *Chromatography* <621>, *System Suitability*.)

Mode: LC

Detector: UV 280 nm

Column: 4.6-mm \times 5.0-cm; 5- μ m packing L1

Column temperature: 40°

Flow rate: 1.2 mL/min

Injection volume: 10 μ L

Run time: NLT 2.0 times the retention time of quetiapine

System suitability

Sample: *Standard solution*

Suitability requirements

Tailing factor: NMT 2.0

Relative standard deviation: NMT 3.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the concentration, C_i , of quetiapine ($C_{21}H_{25}N_3O_2S$) in *Medium* (mg/mL) after each time point (i):

$$C_i = (r_U/r_S) \times C_S \times (M_{r1}/M_{r2}) \times N$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of USP Quetiapine Fumarate RS in the *Standard solution* (mg/mL)

M_{r1} = molecular weight of quetiapine free base, 383.51

M_{r2} = molecular weight of quetiapine fumarate, 883.09

N = number of moles of quetiapine free base per mole of quetiapine fumarate, 2

Calculate the percentage of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at each time point (i):

$$\text{Result}_1 = C_i \times V \times (1/L) \times 100$$

$$\text{Result}_2 = \{[C_2 \times (V - V_3)] + (C_1 \times V_3)\} \times (1/L) \times 100$$

$$\text{Result}_3 = \left\{ \frac{C_3 \times [V - (2 \times V_3)]}{100} + [(C_2 + C_1) \times V_3] \right\} \times (1/L) \times 100$$

$$\text{Result}_4 = \left\{ \frac{C_4 \times [V - (3 \times V_3)]}{100} + [(C_3 + C_2 + C_1) \times V_3] \right\} \times (1/L) \times 100$$

C_i = concentration of quetiapine in *Medium* in the portion of sample withdrawn at each time point (mg/mL)

V = volume of *Medium*, 900 mL

L = label claim (mg/Tablet)

V_3 = volume of the *Sample solution* withdrawn from the *Medium* (mL)

Tolerances: See *Table 6*.

Table 6

Time Point (i)	Time (h)	Amount Dissolved (for 50- and 150-mg Tablets) (%)	Amount Dissolved (for 200-, 300-, and 400-mg Tablets) (%)
1	1	NMT 15	NMT 10
2	4	21–41	21–41
3	8	56–76	51–71
4	16	NLT 80	NLT 80

The percentages of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

Test 7: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 7*.

Acid stage medium: Citrate buffer, pH 4.8 (9.6 g/L of anhydrous citric acid in water prepared as follows.

Transfer a suitable quantity of anhydrous citric acid to an appropriate volumetric flask. Dissolve in 60% of the flask volume of water, then add 9% of the flask volume of 1 N sodium hydroxide VS. Dilute with water to volume); 900 mL, deaerated

0.05 M phosphate buffer solution: 17.9 g/L of dibasic sodium phosphate dodecahydrate solution prepared as follows. Transfer a suitable amount of dibasic sodium phosphate dodecahydrate to an appropriate volumetric flask containing 40% of the flask volume of water. Add 46% of the flask volume of 1 N sodium hydroxide VS and dilute with water to volume.

Buffer stage medium: Phosphate buffer, pH 6.6 (add 100 mL of 0.05 M phosphate buffer solution to the *Acid stage medium*; adjust with 1 N sodium hydroxide VS or 1 N hydrochloric acid VS to obtain a pH of 6.6 ± 0.20 , if necessary); 1000 mL

Apparatus 1: 20-mesh basket; 200 rpm

Times: 1 and 4 h in *Acid stage medium*; 6, 10, and 16 h in *Buffer stage medium*. The time in the *Buffer stage medium* includes the time in the *Acid stage medium*.

Procedure: Run the test in the *Acid stage medium* for the times specified. After 5 h, add 100 mL of 0.05 M phosphate buffer solution and continue running the test in *Buffer stage medium* for the times specified.

Standard stock solution: 1.2 mg/mL of USP Quetiapine Fumarate RS prepared as follows. Transfer an appropriate amount of USP Quetiapine Fumarate RS into a suitable volumetric flask. Add 40% of the flask volume of methanol and sonicate to dissolve. Dilute with water to volume.

Acid stage standard solution: 0.03 mg/mL of USP Quetiapine Fumarate RS from *Standard stock solution* in *Acid stage medium*

Buffer stage standard solution: 0.03 mg/mL of USP Quetiapine Fumarate RS from *Standard stock solution* in *Buffer stage medium*

Acid stage sample solution: Pass a portion of the solution under test through a suitable filter, discarding the first portion of filtrate if needed. Dilute the filtrate further with *Acid stage medium*, if needed. Replace the portion of solution removed from the vessel with an equivalent volume of warmed *Acid stage medium*.

Buffer stage sample solution: Pass a portion of the solution under test through a suitable filter, discarding the first portion of filtrate if needed. Dilute the filtrate further with *Buffer stage medium*, if needed. Replace the portion of solution removed from the vessel with an equivalent volume of warmed *Buffer stage medium*.

Instrumental conditions

(See *Ultraviolet-Visible Spectroscopy* (857).)

Mode: UV

Analytical wavelength: 290 nm

Blank: *Acid stage medium* or *Buffer stage medium*

System suitability

Samples: *Acid stage standard solution* and *Buffer stage standard solution*

Suitability requirements

Relative standard deviation: NMT 2.0%, *Acid stage standard solution* and *Buffer stage standard solution*

Analysis

Samples: *Acid stage standard solution* and *Acid stage sample solution* or *Buffer stage standard solution* and *Buffer stage sample solution*

Calculate the concentration (C_i) of quetiapine ($C_{21}H_{25}N_3O_2S$) in the sample withdrawn from the vessel at each time point (i):

$$\text{Result}_i = (A_U/A_S) \times C_S \times D \times (M_{r1}/M_{r2}) \times N$$

A_U = absorbance of the *Acid stage sample solution* or *Buffer stage sample solution*

A_S = absorbance of the *Acid stage standard solution* or *Buffer stage standard solution*, corresponding to the related sample solution

C_S = concentration of quetiapine fumarate in *Acid stage standard solution* or *Buffer stage standard solution* (mg/mL)

D = dilution factor for the *Acid stage sample solution* or *Buffer stage sample solution*, if needed

M_{r1} = molecular weight of quetiapine free base, 383.51

M_{r2} = molecular weight of quetiapine fumarate, 883.09

N = number of moles of quetiapine free base per mole of quetiapine fumarate, 2

Calculate the percentage of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at each time point (i):

$$\begin{aligned} \text{Result}_1 &= C_1 \times V_A \times (1/L) \times 100 \\ \text{Result}_2 &= [(C_2 \times V_A) + (C_1 \times V_S)] \times (1/L) \times 100 \\ \text{Result}_3 &= \{(C_3 \times V_B) + [(C_2 + C_1) \times V_S]\} \times (1/L) \times 100 \\ \text{Result}_4 &= \{(C_4 \times V_B) + [(C_3 + C_2 + C_1) \times V_S]\} \times (1/L) \times 100 \\ \text{Result}_5 &= \{(C_5 \times V_B) + [(C_4 + C_3 + C_2 + C_1) \times V_S]\} \times (1/L) \times 100 \end{aligned}$$

C_i = concentration of quetiapine in *Acid stage medium* or *Buffer stage medium* in the portion of sample withdrawn at each time point (mg/mL)

V_A = volume of *Acid stage medium*, 900 mL

L = label claim (mg/Tablet)

V_S = volume of the *Acid stage sample solution* or *Buffer stage sample solution* withdrawn from the vessel and replaced with *Acid stage medium* or *Buffer stage medium*, respectively (mL)

V_B = volume of *Buffer stage medium*, 1000 mL

Tolerances: See *Table 7*.

Table 7

Time Point (i)	Time (h)	Amount Dissolved (%)
1	1	NMT 15
2	4	28–48
3	6	40–60
4	10	62–82
5	16	NLT 80

The percentages of the labeled amount of quetiapine ($C_{21}H_{25}N_3O_2S$) dissolved at the times specified conform to *Dissolution* (711), *Acceptance Table 2*.▲ (RB 1-Apr-2019)

• **UNIFORMITY OF DOSAGE UNITS** (905): Meet the requirements

IMPURITIES

Change to read:

• **ORGANIC IMPURITIES**

Buffer, Mobile phase, Diluent, System suitability solution, Sample solution, and Chromatographic system: Proceed as directed in the *Assay*.

System suitability

Sample: *System suitability solution*

[NOTE—See ▲ *Table 8*▲ (RB 1-Apr-2019) for the relative retention times.]

Suitability requirements

Resolution: NLT 1.5 between quetiapine related compound G and quetiapine related compound H; NLT 2.0 between the quetiapine desthoxy and quetiapine peaks

Analysis

Sample: *Sample solution*

[NOTE—See ▲ *Table 8*▲ (RB 1-Apr-2019) for the relative retention times.]

Calculate the percentage of each degradation product in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (1/F) \times 100$$

r_U = peak response of each degradation product from the *Sample solution*

r_S = peak response of quetiapine from the *Sample solution*

F = relative response factor for the corresponding degradation product from ▲ *Table 8*▲ (RB 1-Apr-2019)

Acceptance criteria: See ▲ *Table 8*.▲ (RB 1-Apr-2019) Disregard peaks less than 0.05%.

▲Table 8▲ (RB 1-Apr-2019)

Name	Relative Retention Time	Relative Response Factor	Acceptance Criteria, NMT (%)
Fumaric acid ^a	0.1	—	—
Quetiapine related compound G	0.48	1.4	0.2
Quetiapine related compound H	0.57	1.0	0.2
Quetiapine desethoxy ^b	0.87	—	—
Quetiapine	1.0	—	—
Quetiapine related compound B ^b	1.9	—	—
Any individual unspecified degradation product	—	1.0	0.2
Total degradation products	—	—	0.4

^a Counter ion peak, not to be included in the total degradation products.

^b Process impurity controlled in the drug substance. Included for identification purposes only. Not reported for the drug product and not included in the total degradation products.

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in well-closed containers. Store at controlled room temperature.
- **LABELING:** When more than one *Dissolution* test is given, the labeling states the *Dissolution* test used only if *Test 1* is not used.
- **USP REFERENCE STANDARDS** <11>
 - USP Quetiapine Fumarate RS
 - USP Quetiapine Related Compound H RS
4-(Dibenzo[*b,f*][1,4]thiazepin-11-yl)-1-[2-(2-hydroxyethoxy)ethyl]piperazine 1-oxide.
C₂₁H₂₅N₃O₃S 399.51
 - USP Quetiapine System Suitability RS
It contains quetiapine fumarate and at least 0.1% of each of the following impurities:
Quetiapine related compound B: 11-(Piperazin-1-yl)dibenzo[*b,f*][1,4]thiazepine; Quetiapine related compound G: Dibenzo[*b,f*][1,4]thiazepin-11(10*H*)-one; and Quetiapine desethoxy: 2-[4-(Dibenzo[*b,f*][1,4]thiazepin-11-yl)piperazin-1-yl]ethanol.