

[x](#)

How to Use

- **Searching:** Type keyword in search field at top of page. Search by all or part of a monograph title. For searches using multiple criteria, you will find items that match each of the specified criteria unless quotation marks are used.
 - For example, a search on Aminosalicyclic Acid Tablets will result in anything that contains “Aminosalicyclic” OR “Acid” OR “Tablets”
 - A search for “Aminosalicyclic Acid Tablets” will result in anything that specifically contains “Aminosalicyclic Acid Tablets”
- **Sorting:** Click on any column header title to sort alphabetically or chronologically in ascending or descending order. Note: the page load column is sorted alphabetically so that a number is ordered by first digit vs. by the actual number; thus, numbers will not always be in order.
 - For example, page 2178 will come before page 74 on a page sort.
- **Downloading:** You can download the Errata table in Comma-separated Value (.csv). The download will include the Errata that you have filtered on.
- **Importing:** You will need to import the file into Excel or Open Office with UTF-8 encoding, as opposed to simply opening it. To import, open Excel or Open Office and select import from the File drop-down. Depending on the version you are using, you should be presented with import formatting options to include UTF-8 as one of the first steps. Importing via UTF-8 should eliminate odd character conversions.

| Monograph Title | Section | Source | Page Number | Errata Post | Errata Official | Target Errata | Target Online | Description |
|---------------------------------|-------------------------|-----------------------------|-----------------------------|-----------------------------|---------------------------------|-----------------------------------|---------------------------------|-------------|
| | | Publication | | Date | Date | Print Publication | Fix Publication | |
| THIMEROSAL | SPECIFIC | USP36–NF31 | 5369 | 31-May-2013 | 1-Jun-2013 | USP37–NF32 | USP37–NF32 | Line 4 of |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|--|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|---|
| TOPICAL AEROSOL | TESTS/ <i>Alcohol Determination, Method II</i> | | | | | | | | <p><i>Analysis:</i> Change Determine the alcohol content of the sample thus prepared by the <i>Gas-Liquid Chromatographic Method (see Method II in Alcohol Determination <611>, using methyl ethyl ketone as the internal standard in place of acetone.</i></p> <p>to: Determine the alcohol content of the sample thus prepared by the <i>Gas Chromatographic Method (see Method II in Alcohol D</i></p> |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|------------------------------------|-------------------------------|---------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|---|
| CARAWAY OIL | DEFINITION | USP36–NF31 | 1924 | 31-May-2013 | | 1-Jun-2013 | USP37–NF32 | USP37–NF32 | <p><i>ete rminat ion <611></i>), using methyl ethyl ketone as the internal standard in place of acetonitrile.</p> <p>Line 3: Change It contains NMT 50.0% of <i>d</i>-carvone (C₁₀H₁₄O). to: It contains NLT 50.0% of <i>d</i>-carvone (C₁₀H₁₄O).</p> |
| NORTRIPTYLIN NE HYDROCHLORIDE | IMPURITIES/Organic Impurities | <i>First Supplement to USP36–NF31</i> | 6027 | 31-May-2013 | | 1-Jun-2013 | USP37–NF32 | USP37–NF32 | <p>Line 3 of <i>Acceptance criteria</i>: Change <i>Standard solution</i> to: <i>Sample solution</i></p> |
| FOSPHENYTOIN N SODIUM INJECTION | USP Reference standards <11> | USP36–NF31 | 3680 | 31-May-2013 | | 1-Jun-2013 | USP37–NF32 | USP37–NF32 | <p>Line 6: Change C₁₄H₁₅NO₂ to: C₁₄H₁₃NO₂</p> |
| MINOCYCLINE | Assay | USP36–NF31 | 4378 | 31-May-2013 | | 1-Jun-2013 | USP37–NF32 | USP37–NF32 | <p>Line 2: Change</p> |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|---|
| HYDROCHLORIDE TABLETS | | | | | | | <p><i>Mobile phase, Standard preparation, Resolution solution, and Chromatographic system</i></p> <p>—Proceed as directed in the Assay under <i>Minocycline Hydrochloride</i>.</p> <p>to:</p> <p><i>Mobile phase</i>—Prepare a mixture of 0.2 M ammonium oxalate, 0.01 M edetate disodium, dimethylformamide, and tetrahydrofuran (600:180:120:80). Adjust with ammonium hydroxide to a pH of 7.2, and pass through a filter of 0.5-μm</p> |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|---|
| | | | | | | | <p>or finer pore size. Make adjustments if necessary (see <i>System Suitability</i> under <i>Chromatography</i> <621>). <i>Standard preparation</i></p> <p>—Dissolve an accurately weighed quantity of USP Minocycline Hydrochloride RS in water to obtain a solution having a known concentration of about 500 µg of minocycline (C₂₃H₂₇N₃O₇) per mL. Use this solution within 3 hours. <i>Resolution solution</i></p> <p>—Transfer 10</p> |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|--|
| | | | | | | | <p>mg of USP Minocycline Hydrochloride RS to a 25-mL volumetric flask, add 20 mL of 0.2 M ammonium oxalate, and swirl to dissolve. Heat on a water bath at 60° for 180 minutes, and allow to cool. Dilute with water to volume, and mix.</p> <p><i>Chromatographic system (see Chromatography <621>)</i>—The liquid chromatograph is equipped with a 280-nm detector and a 4.6-mm x 25-cm column that contains</p> |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|--|
| | | | | | | | <p>5-μm packing L1, and is maintained at a constant temperature of about 40°. The flow rate is about 1.5 mL per minute. Chromatograph the <i>Standard preparation</i>, and record the peak responses as directed for <i>Procedure</i>: the capacity factor, k', is not less than 5.0 and not more than 11.5; the tailing factor for the analyte peak is not less than 0.9 and not more than 2.0; and the relative standard deviation for replicate injections is not</p> |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|---|
| | | | | | | | <p>more than 2.0%. Chromatograph the <i>Resolution solution</i>, and record the peak responses as directed for <i>Procedure</i>: the relative retention times are about 0.7 for epiminocycline and 1.0 for minocycline; and the resolution, <i>R</i>, between epiminocycline and minocycline is not less than 4.6.</p> <p>AND</p> <p>Line 1 of <i>Procedure</i>: Change Proceed as directed for <i>Procedure</i> in the Assay under</p> |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|--|---|-----------------------------|---|---|--|--|---|
| | | | | | | | <p><i>Minocycline Hydrochloride.</i> to: Separately inject equal volumes (about 20 µL) of the <i>Standard preparation</i> and the <i>Assay preparation</i> into the chromatograph, record the chromatograms, and measure the responses for the major peaks.</p> <p>Line 3 of <i>System suitability.</i> Change [Note—The retention time for squalene is about 18 min; the relative retention times for tetradecane, hexadecane,</p> |
| HYDROGENAT ASSAY/ ED POLYDECENE | <i>Content of Decene Oligomer</i> USP36–NF31 | 2133 | 29-Mar-2013 | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|--|
| CLOTRIMAZOL ASSAY/ E AND BETAM ETHASONE DI PROPIONATE CREAM | USP36–NF31 | 3075 | 29-Mar-2013 | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | and squalene are about 0.5, 0.6, and 1.0, respectively.] to: [Note—The retention time for squalene is about 18 min; the relative retention times for tetradecane, hexadecane, and squalene are about 0.5, 0.6, and 1.0, respectively.] Line 3 of <i>Betamethasone dipropionate stock solution</i> : Change <i>J</i> being the ratio (in mg/g) of betamethasone to clotrimazole in the Cream to: <i>J</i> being the ratio of the labeled amount of |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|---|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|--|
| CALCIUM SULFATE | SPECIFIC TESTS/ <i>Loss on Drying</i> <731> | USP35–NF30 | 1724 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | betamethasone (in mg/g) to the labeled amount of clotrimazole (in mg/g) in the Cream Line 1 of <i>Acceptance criteria</i> : Change NMT 1.5% for the anhydrous form and NMT 19.0%–23.0% for the dihydrate to: NMT 1.5% for the anhydrous form and 19.0%–23.0% for the dihydrate |
| TAPIOCA STARCH | <i>Limit of oxidizing substances</i> | USP35–NF30 | 1987 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Line 8: Change Add 1 mL of starch TS, and titrate with 0.002 N sodium thiosulfate VS to the disappearance of the starch–iodide color. |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|-------------------------------------|---|--|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|---|
| LEVETIRACETAM | ADDITIONAL REQUIREMENTS | USP35–NF30 | 3659 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | to: Add 1 mL of starch TS, and titrate with 0.002 N sodium thiosulfate VS to the disappearance of the starch–iodine color. Line 9 of <i>USP Reference Standards</i> <11>: Change $C_8H_{14}ClNO_3$ 207.65 to: $C_8H_{15}ClN_2O_2$ 206.67 |
| DULOXETINE DELAYED-RELEASE CAPSULES | PERFORMANCE TESTS/ <i>Dissolution <711>/ Chromatographic system</i> | <i>Second Supplement to USP35–NF30</i> | 5940 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Line 1 of <i>Column</i> : Change 4.6-mm x 7.5-cm; 3- μ m packing L7 to: 4.6-mm x 7.5-cm; 3- or 3.5- μ m packing L7 |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|---|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|---|
| HYMETELLOS E | IM PUR ITIES/ <i>Chloride and Sulfate, Chloride <221></i> | USP36–NF31 | 2044 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Change the subsection title <i>Standard solution</i> to: <i>Control solution AND Line 4 of Analysis: Change Standard solution</i> to: <i>Control solution AND Line 2 of Acceptance criteria: Change Standard solution</i> to: <i>Control solution</i> |
| BENZTROPINE MESYLATE | CHEMICAL INFORMATION | USP36–NF31 | 2628 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Line 2: Change 8-Azabicyclo[3.2.1]octane, 3-(di phenylmethoxy) -, <i>endo</i> -, methanesulfonate; to: 8-Azabicyclo[3.2.1]octane, 3-(di |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|---|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|--|
| LORAZEPAM TABLETS | IMPURITIES/Organic Impurities/System suitability/Suitability requirements | USP36–NF31 | 4153 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | phenylmethoxy)-N-methyl-, endo-, methane sulfonate; Line 1 of <i>Tailing factor</i> . Change 2.0, <i>Standard solution</i> to: NMT 2.0, <i>Standard solution</i> |
| BRINZOLAMIDE | Related compounds/Test 2 | USP35–NF30 | 2385 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Line 15 of <i>Procedure</i> : Change relative retention time greater than 6. to: relative retention greater than 6. |
| VINORELBINE INJECTION | Assay | USP35–NF30 | 5028 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Line 1: Change <i>Phosphate buffer</i> , <i>Mobile phase</i> , and <i>System suitability solution</i> —Proceed as directed in the |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|---|
| | | | | | | | <p>Assay under <i>Vinorelbine Tartrate</i>. to: <i>Phosphate buffer</i>—Dissolve 6.9 g of monobasic sodium phosphate in 900 mL of water. Adjust with phosphoric acid to a pH of 4.2, dilute with water to 1000 mL, and mix.</p> <p><i>Mobile phase</i>—Dissolve 1.22 g of sodium 1-decanesulfonate in 620 mL of methanol. Add 380 mL of <i>Phosphate buffer</i>, mix, filter, and degas. Make adjustments if necessary (see</p> |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|---|
| | | | | | | | <p><i>System Suitability</i> under <i>Chromatography</i> <621>).</p> <p><i>System suitability solution</i></p> <p>—Dissolve accurately weighed quantities of USP Vinorelbine Tartrate RS and USP Vinorelbine Related Compound A RS in water, and dilute quantitatively, and stepwise if necessary, with water to obtain a solution having known concentrations of about 1.4 mg per mL and 0.01 mg per mL,</p> |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|--|
| POWDERED BLACK PEPPER EXTRACT | DEFINITION | USP36–NF31 1365 | 29-Mar-2013 | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | <p>respectively. Expose a portion of this solution in a suitable xenon lamp apparatus capable of supplying a dose of 1600 KJ/m² between 310 and 800 nm at a power of 500 W/m² for about 1 h, in order to generate an additional degradation product 3,6-epoxy vinorelbine having a relative retention time of about 0.8.</p> <p>Line 5: Change It contains NLT 90.0% and NMT 110.0% of the labeled amount of piperine.</p> |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|--|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|--|
| POLYVINYL ACETATE PHTHALATE | IM PURITIES/ <i>Free Phthalic Acid</i> | USP36–NF31 | 2168 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | to: It contains NLT 90.0% and NMT 110.0% of the labeled amount of piperine, calculated on the dried basis. Line 1 of <i>Sample solution</i> : Change 6 mg/mL of polyvinyl acetate to: 6 mg/mL of polyvinyl acetate phthalate |
| DILTIAZEM HYDROCHLORIDE ORAL SUSPENSION | ASSAY/ <i>Procedure</i> | USP36–NF31 | 3263 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Line 6 of <i>Sample solution</i> : Change Pipet 1.0 mL of the sample solution to: Pipet 1.0 mL of the sample |
| IFOSFAMIDE | <i>Chloroform-</i> | USP35–NF30 | 3477 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Line 18 of <i>Test</i> |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|---|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|---|
| | | | | | | | | | <i>preparation:</i> Change ammonium hydroxide solution. to: ammonium hydroxide. |
| ATROPINE SULFATE TABLETS | Assay | USP35–NF30 | 2272 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Line 9 of <i>Procedure:</i> Change R_U and R_S are as defined therein. to: R_U and R_S are the peak area ratios of atropine to homatropine. |
| VANCOMYCIN INJECTION | SPECIFIC TESTS/ <i>Composition of Vancomycin</i> | USP35–NF30 | 5003 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Line 16 of <i>Analysis:</i> Change D = dilution factor, <i>Sample stock solution</i> to <i>Sample solution</i> , 25 to: D = dilution factor, <i>Sample</i> |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|---|--|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|---|
| | | | | | | | | | <p><i>stock solution to Sample solution AND</i></p> <p>Line 29 of <i>Analysis:</i></p> <p>Change <i>D</i> = dilution factor, <i>Sample stock solution to Sample solution</i>, 25 to:</p> <p><i>D</i> = dilution factor, <i>Sample stock solution to Sample solution</i></p> |
| SUMATRIPTAN INJECTION | SPECIFIC TESTS/ <i>Osmolality and Osmolarity <785></i> | <i>Second Supplement to USP35–NF30</i> | 5996 | 29-Mar-2013 | | 1-Apr-2013 | <i>USP37–NF32</i> | <i>USP37–NF32</i> | <p>Line 1: Change 270–330 mOsmol to:</p> <p>270–330 mOsmol/kg</p> |
| INOSITOL | SPECIFIC TESTS/ <i>Conductivity</i> | <i>USP36–NF31</i> | 2049 | 29-Mar-2013 | | 1-Apr-2013 | <i>USP37–NF32</i> | <i>USP37–NF32</i> | <p>Line 1 of <i>Sample solution:</i></p> <p>Change Transfer 10.0 g of Inositol, weighed and calculated on the dried basis,</p> |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|--|------------------------------------|-----------------------------|---|--------------------------------------|---|---|---|
| CLONAZEPAM ASSAY/ ORAL <i>Procedure</i> SUSPENSION | USP36–NF31 | 3053 | 29-Mar-2013 | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | to a 50-mL volumetric flask, and dissolve in and dilute with water (previously boiled and cooled to room temperature) to volume. to: 0.2 g/mL of Inositol in water (previously boiled and cooled to room temperature). Line 6 of <i>Sample solution</i> : Change Pipet 2.5 mL of the <i>Sample solution</i> to: Pipet 2.5 mL of the sample |
| METHENAMIN ASSAY/ E ORAL <i>Procedure</i> SOLUTION | USP36–NF31 | 4288 | 29-Mar-2013 | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Line 10 of <i>Analysis</i> : Change <i>B</i> |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|--|---|-----------------------------|---|--|---|---|--|
| AZITHROMYCI N | IM PUR ITIES/ <i>Organic I mpurities/Proce dure 2</i> | USP35–NF30 2279 | 29-Mar-2013 | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | <p>$S =$ absorbance of the <i>Sample blank</i> to: $B_S =$ absorbance of the <i>Standard blank</i></p> <p>Line 15 of <i>Analysis:</i> Change $C_S =$ concentration of USP Azithromycin RS in the <i>Standard solution</i> ($\mu\text{g/mL}$) to: $C_S =$ concentration of USP Azithromycin RS in the <i>Standard solution</i> (mg/mL) AND Add after C_U: $P =$ potency of USP</p> |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Errata Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|-------------------------------------|--|--|-----------------------------|----------------------------------|---------------------------------------|--------------------------------------|---|---|--|
| ATRACURIUM BESYLATE INJECTION | IM PUR ITIES/ <i>Organic Impurities/Acceptance criteria/</i> Table 2 | <i>Second Supplement to USP35–NF30</i> | 5909 | 29-Mar-2013 | | 1-Apr-2013 | <i>USP37–NF32</i> | <i>USP37–NF32</i> | Azithromycin RS (µg/mg of azithromycin) Footnote b: Change <i>cis</i> isomer of the hydroxy compound. to: <i>trans</i> isomer of the hydroxy compound. AND Footnote c: Change <i>trans</i> isomer of the hydroxy compound. to: <i>cis</i> isomer of the hydroxy compound. AND Footnote d: Change <i>cis</i> isomer of the monoacrylate. to: <i>trans</i> isomer of the |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|--|-----------------------------|---|--------------------------------------|---|---|--|
| ZINC SULFATE TABLETS | Identification/B. <i>Zinc</i> <i>USP35–NF30</i> | 5077 | 29-Mar-2013 | 1-Apr-2013 | <i>USP37–NF32</i> | <i>USP37–NF32</i> | monoacrylate. AND Footnote e: Change <i>trans</i> isomer of the monoacrylate. to: <i>cis</i> isomer of the monoacrylate. Line 1 of <i>Sodium hydroxide solution:</i> Change 42 mg/mL of sodium hydroxide to: 420 mg/mL of sodium hydroxide |
| DIETHYL SEBACATE | DEFINITION <i>USP36–NF31</i> | 1994 | 29-Mar-2013 | 1-Apr-2013 | <i>USP37–NF32</i> | <i>USP37–NF32</i> | Line 2: Change Diethyl Sebacate consists of the diester of alcohol and sebacic acid. to: |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|---|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|---|
| PROPYLENE GLYCOL MONOLAURATE | IM PURITIES/ <i>Limit of Propylene Glycol</i> | USP36–NF31 | 2180 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Diethyl Sebacate consists of the diester of alcohol (ethanol) and sebacic acid. Line 8 of <i>Analysis</i> : Change Calculate the percentage of free propylene glycol in the portion of Propylene Glycol Monocaprylate taken: to: Calculate the percentage of free propylene glycol in the portion of Propylene Glycol Monolaurate taken: Line 5 of <i>Analysis</i> : |
| GLUCONOLAC TONE | IDENTIFICATIO N/A. | USP36–NF31 | 3742 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|--|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|---|
| METHOTREXATE | IMPURITIES/ <i>Organic Impurities/Procedure 1: Related Compounds</i> | USP35–NF30 | 3855 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Change crystals of the phenylhydrazine of gluconic acid to: crystals of the phenylhydrazide of gluconic acid Footnote b of <i>Impurity Table 1</i> : Change (S)-2-{4-[(2-Amino-4-oxo-1,4-dihydropteridin-6-yl)methylamino]-N-methylbenzamide}pentanedioic acid. to: (S)-2-(4-[(2-Amino-4-oxo-1,4-dihydropteridin-6-yl)methylamino]benzamide)pentanedioic acid. |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|-------------------------------------|-----------------------------|---|--------------------------------------|---|---|--|
| BETAMETHAS ONE ORAL SOLUTION | <i>Identification/A:</i> USP35–NF30 | 2336 | 29-Mar-2013 | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | acid. Line 1: Change A: to: <i>A: Thin-Layer Chromatographic Identification Test <201></i> — |
| VINORELBINE INJECTION | <i>Related compounds</i> USP35–NF30 | 5028 | 29-Mar-2013 | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Delete the subsection <i>Standard solution</i> and <i>Diluted standard solution</i> . Replace with: <i>Standard solution</i> —Dissolve an accurately weighed quantity of USP Vinorelbine Tartrate RS in <i>Mobile phase</i> to obtain a solution having a known concentration of about 1.4 mg per mL. |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|--|
| | | | | | | | <p><i>Diluted standard solution</i> —Transfer 1.0 mL of the <i>Standard solution</i> to a 50-mL volumetric flask, and dilute with <i>Mobile phase</i> to volume. Pipet 1.0 mL of this solution into a 100-mL volumetric flask, and dilute with <i>Mobile phase</i> to volume.</p> <p>AND</p> <p>Line 1 of <i>Procedure</i>: Change Proceed as directed for <i>Procedure</i> in the test for <i>Related compounds</i> under <i>Vinorelbine</i></p> |


| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|--|
| | | | | | | | <p><i>Tartrate.</i></p> <p>to:</p> <p>Separately inject equal volumes (about 20 µL) of the <i>Test solution</i> and the <i>Diluted standard solution</i> into the chromatograph, record the chromatograms, and measure the areas for the major peaks. Record the chromatograms for three times the retention time of the vinorelbine peak. Disregard any peaks with an area less than or equal to one-half of the area of the peak obtained for vinorelbine in</p> |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|--|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|--|
| | | | | | | | | | <p>the <i>Diluted standard solution</i>. Calculate the percentage of each impurity in the portion of Injection taken by the formula: $100(r_i/r_s)$ in which r_i is the peak response for each impurity obtained from the <i>Test solution</i>; and r_s is the sum of the responses of all the peaks.</p> |
| GOOD STORAGE AND DISTRIBUTION PRACTICES FOR DRUG PRODUCTS | QUALITY MANAGEMENT SYSTEM/Storage Management System/Receiving and Transferring Drug Products | USP36–NF31 | 693 | 29-Mar-2013 | | 1-Apr-2013 | USP37–NF32 | USP37–NF32 | Line 1 of footnote 1: Change JP Edmond, to: JP Emond, |
| ACESULFAME POTASSIUM | IMPURITIES/Limit | USP35–NF30 | 1680 | 31-Jan-2013 | | 1-Feb-2013 | USP37–NF32 | Second Supplement to | Line 25 of Analysis: |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Errata Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|-----------------------------------|--|------------------------------------|-----------------------------|----------------------------------|---------------------------------------|--------------------------------------|---|---|--|
| | | <i>of Fluoride</i> | | | | | | USP36–NF31 | Change C = concentration of fluoride in the <i>Sample solution</i> , from the standard curve (mg/mL) to: C = concentration of fluoride in the <i>Sample solution</i> , from the standard curve (µg/mL) |
| AMANTADINE HYDROCHLORIDE CAPSULES | PERFORMANCE TESTS/ <i>Dissolution</i> <711>/Test 2 | USP35–NF30 | 2153 | 31-Jan-2013 | | 1-Feb-2013 | USP37–NF32 | Second Supplement to USP36–NF31 | Line 5 of <i>Chromatographic system</i> : Change Column: 0.32-mm x 30-cm, 0.25-µm film, phase G1 to: Column: 0.32-mm x 30-m, 0.25-µm film, phase G1 |
| OXYBUTYNIN CHLORIDE EX E | PERFORMANCE TESTS/ <i>Dissolution</i> <711>/Test 2 | USP35–NF30 | 4167 | 31-Jan-2013 | | 1-Feb-2013 | USP37–NF32 | Second Supplement to | Line 3 of <i>Working</i> |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|--|---------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|--|
| TENDED-RELEASE TABLETS | TESTS/ <i>Dissolution</i> <711>/Test 2 | | | | | | | USP36–NF31 | <i>standard solution:</i> Change or transfer 10 mL for Tablets labeled to contain 10 mg, to a 100-mL volumetric flask. to: transfer 10 mL for Tablets labeled to contain 10 mg, or transfer 15 mL for Tablets labeled to contain 15 mg to a 100-mL volumetric flask. |
| ALLANTOIN | IDENTIFICATION | <i>First Supplement to USP35–NF30</i> | 5429 | 31-Jan-2013 | | 1-Feb-2013 | USP37–NF32 | <i>Second Supplement to USP36–NF31</i> | Line 1 of <i>B. Thin-Layer Chromatographic Identification Test <201></i> : Change The R_F value of the principal spot from <i>Sample solution A</i> corresponds |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|-------------------------|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|--|
| CAPTOPRIL ORAL SUSPENSION | Assay | USP35–NF30 | 2477 | 31-Jan-2013 | | 1-Feb-2013 | USP37–NF32 | Second Supplement to USP36–NF31 | <p>to that from <i>Standard solution A</i>, as described in the test for <i>Organic Impurities</i>.</p> <p>to:</p> <p>The R_F value of the principal spot from <i>Sample solution B</i> corresponds to that from <i>Standard solution A</i>, as described in the test for <i>Organic Impurities</i>.</p> <p>Line 1 of <i>Mobile phase</i>: Change Prepare a filtered and degassed mixture of methanol and water (11:9) containing 0.5 mL of phosphoric acid.</p> <p>to:</p> |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|---------------------------------------|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|---|
| CHROMATOGRAPHY | SYSTEMS UNIT ABILITY/Stationary Phase | USP35–NF30 | 258 | 31-Jan-2013 | | 1-Feb-2013 | USP37–NF32 | Second Supplement to USP36–NF31 | <p>Methanol and water (55:45) containing 0.5 mL/L of phosphoric acid. Filter, and degas.</p> <p>Line 3 of <i>Flow Rate (HPLC)</i>: Change </p> <p>in which F_1 is the flow rate indicated in the monograph, in mL/min; F_2 is the adjusted flow rate, in mL/min; l_1 is the length of the column indicated in the monograph; l_2 is the length of the column used; d_1 is the column inner diameter indicated in the monograph; and d_2 is the</p> |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|--|
| LOW-SUBSTIT Assay | USP35–NF30 | 1822 | 31-Jan-2013 | 1-Feb-2013 | USP37–NF32 | Second | <p>internal diameter of the column used. Additionally, the flow rate can be adjusted by $\pm 50\%$.</p> <p>to:</p> <p>⊗</p> <p>in which F_1 is the flow rate indicated in the monograph, in mL/min; F_2 is the adjusted flow rate, in mL/min; d_1 is the column inner diameter indicated in the monograph; and d_2 is the internal diameter of the column used. Additionally, the flow rate can be adjusted by $\pm 50\%$.</p> <p>Line 2: Change</p> |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|---------------------------------------|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|--|
| UTED HYDROXYPROPYL CELLULOSE | | | | | | | | <i>Supplement to USP36–NF31</i> | <i>Hypromellose 2906, except to substitute Low-Substituted Hydroxypropyl Cellulose for Hypromellose 2906 throughout. to: Hypromellose, except to substitute Low-Substituted Hydroxypropyl Cellulose for Hypromellose throughout.</i> |
| NORGESTIMATE | <i>Limit of residual solvents</i> | <i>USP35–NF30</i> | 4083 | 31-Jan-2013 | | 1-Feb-2013 | <i>USP37–NF32</i> | <i>Second Supplement to USP36–NF31</i> | Line 1 of <i>Limit of residual solvents</i> : Change to: <i>Limit of residual solvents <467></i> |
| VINCRISTINE SULFATE | IMPURITIES/ <i>Organic Impurities</i> | <i>USP35–NF30</i> | 5022 | 31-Jan-2013 | | 1-Feb-2013 | <i>USP37–NF32</i> | <i>Second Supplement to USP36–NF31</i> | Line 5 of <i>Analysis</i> : Change Result = $[r_{UA}] / (r_{UA} + 25r)$ |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|--|---|-----------------------------|----------------------------------|--------------------------------------|---|---|---|
| | | | | | | | | $\frac{r_{UB}}{r_{UA} + 30r_{UB}} \times 100$ to: Result = $\frac{r_{UA}}{r_{UA} + 30r_{UB}} \times 100$ AND Change line 12 of <i>Analysis</i> : Result = $\frac{r_{UA}}{r_{UA} + 25r_{UB}} \times 100$ to: Result = $\frac{r_{UA}}{r_{UA} + 30r_{UB}} \times 100$ |
| ADAPALENE | IM PUR ITIES/ <i>Residual Solvent: Limit of 2012</i>) <i>Triethylamine</i> | <i>Revision Bulletin (Official December 01,</i> | Online | 31-Jan-2013 | 1-Feb-2013 | <i>USP37–NF32</i> | <i>Second Supplement to USP36–NF31</i> | Line 1 of <i>Diluent</i> : Change Dimethyl sulfoxide and 1 N sodium hydroxide solution (4:1) to: Dimethyl sulfoxide AND Line 1 of <i>Standard solution</i> : Change 3.2 ?g/mL of |

| Monograph Title Section | Source Publication | Page Number | Errata Post Date Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---|------------------------------------|-----------------------------|---|--------------------------------------|---|---|---|
| | | | | | | | <p>USP Triethylamine RS in <i>Diluent</i> to: 4.0 ?g/mL of USP Triethylamine RS in <i>Diluent</i>. Transfer 4.0 mL of this solution to a 20-mL headspace vial, and add 1.0 mL of 1 N NaOH solution. AND Line 1 of <i>Sample solution</i>: Change 40 mg/mL of Adapalene in <i>Diluent</i> to: 50 mg/mL of Adapalene in <i>Diluent</i>. Transfer 4.0 mL of this solution to a 20-mL headspace vial,</p> |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|-------------------------|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|---|
| GLYCERYL BEHENATE | IMPURITIES | USP35–NF30 | 1811 | 31-Jan-2013 | | 1-Feb-2013 | USP37–NF32 | Second Supplement to USP36–NF31 | <p>and add 1.0 mL of 1 N NaOH solution.</p> <p>Line 34 of <i>Content of 1-Monoglycerides/Analysis:</i> Change <i>F</i> = equivalency factor of glyceryl monobehenate, 207.3 mg/mEq to: <i>F</i> = equivalency factor of glyceryl monobehenate, 0.2073 g/mEq AND Line 19 of <i>Limit of Free Glycerin/Analysis:</i> Change <i>F</i> = equivalency factor of glycerin, 23.0 mg/mEq to: <i>F</i> = equivalency</p> |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|-------------------------|------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|--|
| CAPTOPRIL ORAL SOLUTION | Assay | USP35–NF30 | 2477 | 31-Jan-2013 | | 1-Feb-2013 | USP37–NF32 | Second Supplement to USP36–NF31 | factor of glycerin, 0.023 g/mEq Line 1 of <i>Mobile phase</i> : Change Prepare a filtered and degassed mixture of methanol and water (11:9) containing 0.5 mL of phosphoric acid. to: Methanol and water (55:45) containing 0.5 mL/L of phosphoric acid. Filter, and degas. |
| PHENYLALANINE IMPURITIES | | USP35–NF30 | 4296 | 31-Jan-2013 | | 1-Feb-2013 | USP37–NF32 | Second Supplement to USP36–NF31 | Line 1 of <i>Heavy Metals, Method I</i> <231>: Change <i>Method I</i> to: <i>Method II</i> |
| ELEMENTAL I | Drug | Second | 5633 | 31-Jan-2013 | | 1-Feb-2013 | USP37–NF32 | Second | Row 16 of |

| Monograph Title | Section | Source Publication | Page Number | Errata Post Date | Sort ascending | Errata Official Date | Target Errata Print Publication | Target Online Fix Publication | Description |
|---------------------------------|--|--------------------------------------|-----------------------------|----------------------------------|--------------------------------|--------------------------------------|---|---|---|
| MPURITIES--LI MITS | <i>Products/Large Volume Parenterals</i> | <i>Supplement to USP35--NF30</i> | | | | | | <i>Supplement to USP36--NF31</i> | Column 4 of <i>Table 1:</i> Change 70 to: 100 AND Row 16 of Column 5 of <i>Table 1:</i> Change 25 to: 10 |

Pagination

- [First page « First](#)
- [Previous page ‹ Previous](#)
- ...
- [Page 32](#)
- [Page 33](#)
- [Page 34](#)
- [Page 35](#)
- [Page 36](#)
- [Page 37](#)
- [Page 38](#)
- [Page 39](#)
- [Page 40](#)
- [Next page Next ›](#)
- [Last page Last »](#)
